

University of Groningen

The neuroanatomical organization of intrinsic brain activity measured by fMRI activity in the human visual cortex

Gravel Araneda, Nicolas Gaspar

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Gravel Araneda, N. G. (2018). *The neuroanatomical organization of intrinsic brain activity measured by fMRI activity in the human visual cortex*. [Thesis fully internal (DIV), University of Groningen]. Rijksuniversiteit Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Invasive electrophysiological recordings of neuronal activity from the visual cortex of cats and other animals have revealed that spontaneous neuronal activity reflects the underlying neuroanatomical organization [30, 70, 40]. However, for non-invasive and indirect functional magnetic resonance imaging (fMRI) recordings of neuronal activity from visual cortical areas, this relationship between intrinsic activity (often referred to as “resting-state”) and cortical neuroanatomical organization is not immediately obvious [42, 36, 62, 11, 27].

When measured using fMRI, intrinsic fluctuations in blood-oxygen level dependent (BOLD) activity are correlated between distant brain regions that are anatomically connected, such as homologous areas in the two hemispheres [6, 73]. For this reason, resting state (RS) fMRI has been widely used to study whole-brain interactions in health and disease [63, 72]. However, interpreting patterns of RS-fMRI activity at a more local scale (e.g., that of the visual cortex) remains challenging as activity in nearby sites can be correlated as a result of either neuroanatomical connections, or metabolic and vascular relationships [12, 47, 45, 34, 58, 46]. Currently, this uncertainty limits the use of RS-fMRI for characterizing visual cortical activity in health and disease.

A better understanding of the relationships between neuronal activity, brain anatomy and hemodynamics would help to develop RS-fMRI as a valuable tool for fundamental and applied —non-invasive— research in humans. Aiming to contribute to the understanding of these relationships, this thesis focuses on the following question: What can we learn about the neuroanatomical organization of the human visual cortex from RS-fMRI recordings?

With this central question in mind, I hypothesize that RS-fMRI activity in the human visual cortex reflects its underlying neuroanatomical organization. Using advanced anatomical MRI techniques and novel fMRI analysis methods, I will characterize and interpret the spatial and temporal organization of in-

trinsic fluctuations in BOLD activity across striate (V1) and extrastriate visual cortex (V2 and V3). Moreover, I will discuss how these fluctuations may be shaped by neuroanatomical, physiological and vascular factors.

In the following sections, I outline the questions and main findings of each experimental chapter, while in the background section, I describe the main anatomical features of the human visual system, summarize the experimental methodologies used in this thesis and briefly describe the emergence of the field of resting state research. The latter section reviews some of the previous studies that my work builds on and helps to remind that the brain is not only concerned with the demands imposed by the environment but also with internally generated dynamics. As we will see in the experimental chapters of my thesis, the neuroanatomical organization of the visual system forms a complex recurrent network of feedforward, feedback and lateral connections that cannot be fully grasped from the classical “sandwich” perspective (perception-cognition-action) of brain function [21] but perhaps—as I will discuss later (in **chapter 6**)—by a more constructivist and enactive view of brain function that accounts for recurrent neural processing and circular causality [71, 38].

1.1 Outline

This thesis comprises four experimental sections. In **chapter 2** I ask whether visuotopic organization¹ can be derived from 7T RS-fMRI activity. Based on the hypothesis that intrinsic fluctuations in BOLD activity reflect underlying neuroanatomical organization, I show that it is possible to map—based on RS-fMRI recordings—cortico-cortical neuronal interactions between V1, V2 and V3 using connective field modeling [32, 27].

Based on the hypothesis that both intrinsic and stimulus-evoked BOLD fluctuations are anchored by the same neuroanatomical connections, in **chapter 3** I ask whether patterns of synchronized fMRI activity in RS and visual field mapping (VFM) are comparable (at 7T). By examining local patterns of phase covariation, I find synchronization clusters that are similar, regardless of whether they are derived from RS or VFM data. However, in activity obtained during VFM, phase synchronization is spatially more extensive than in RS derived activity, reflecting stimulus driven interactions between local responses. Nevertheless, the resemblance between RS and VFM-derived synchronization clusters suggests that they share a common neuroanatomical origin [28].

Methods to analyse and interpret RS-fMRI activity can become valuable tools for the study of human neuronal activity in vivo, in particular if they can be generalized to different magnetic field strengths. To verify this for connective field modeling and cluster synchronization analysis of RS data, in **chapter 4** I study the feasibility of reproducing the main findings of **chapters 2 & 3** using data acquired with a 3T rather than a 7T scanner. Despite the lower resolution and signal-to-noise ratio of the 3T data, I find that the results obtained with it are in fair agreement to those obtained previously with 7T data.

¹When the image’s spatial relationships formed in the retina are preserved in the visual cortex in the form of visual field maps: nearby neurons respond to nearby locations in the image.

In **chapter 5** I ask whether the propagation of BOLD activity within and between visual field maps relates to structured neuronal activity. To explore this question, I implement a modeling approach aimed at disentangling the contributions of local activity and directed interactions in shaping BOLD activity propagation [26]. Applying this approach to 7T fMRI data reveals changes in cortical excitability and directed interactions in RS and VFM, pointing to a task-dependent reconfiguration of local, feedforward and feedback interactions across the visual system [29].

1.2 Background

In this section, I provide a general overview for those interested in studying the human visual cortex using RS-fMRI. I briefly describe the anatomical organization of the human visual system and provide some background on the field of fMRI and resting state measurements. For more detailed accounts on the neuronal and anatomical organization of the human visual cortex see e.g. [37, 74].

1.2.1 Visual pathways

Light, in the form of structured energy distributions, is transduced by photoreceptor cells in the retina (rods and cones) into electrical membrane potentials [24]. These potentials are then shaped by a recurrent cellular network consisting of bipolar, horizontal, and amacrine cells. Together they form a mosaic of functional subunits that contribute to inhibitory and sensitivity-adjustment mechanisms and communicate with the retinal ganglion cells (RGC). The axonal projections from the RGC exit the eye to form the optic nerve. These fiber projections from the two eyes decussate in the optic chiasm, with fibers from the left and right half of each retina going to the left and right hemispheres, respectively. Optic nerve fibers project primarily to the lateral geniculate nucleus (LGN), which relays to occipital visual cortical areas –the topic of the research in this thesis.

1.2.2 Retinotopic organization of the visual cortex

Shaped by evolutionary and developmental constraints, the anatomical circuitry of the human sensory cortices follows the topographic organization of their corresponding sensory surfaces. For the visual system, the result is that the image's spatial relationships formed in the retina are preserved in many regions of the cortex in the form of visual field maps: nearby neurons respond to nearby locations in the image [37]. These retino-cortical maps are said to be retinotopically organized, yet their actual location and shape is determined by different constraints from those that shape the eye [64, 55]. The topographic organization of the primary visual cortex (V1), located along the calcarine sulcus, accommodates an hemifield representation of the retinal image in which the foveal region is greatly magnified –a phenomenon called cortical magnification [65].

The nearby visual field maps V2 and V3 cluster around V1, sharing parallel eccentricity representations. However, the visual field maps in V2 and V3 are discontinuous. They split into dorsal and ventral parts, with angular representations alternating in visual field sign along the horizontal meridian. As a result, V2 and V3 are organized into two quarter hemifields. Other extrastriate visual cortical areas are organized similarly [74]. The multiplicity of visual field maps in the visual cortex appears to reflect the fact that our perceptual machinery is attuned to stable structural features in our habitat. As neurons process different aspects of the image, cortical circuitry is organized into receptive fields that preserve its spatial organization. As a result, cortical areas subserving different functions still preserve this level of organization [74]. In the experimental chapters that will follow, I relate task-dependent changes in fMRI activity to the retinotopic organization of V1, V2 and V3.

1.2.3 Hierarchical organization of the visual cortex

Traditionally, neuroanatomical connections among visual cortical areas have been thought to follow a primarily parallel-hierarchical bottom-up architecture [19]. In this view, parallel neuronal pathways, originating from anatomically and functionally distinct cell types in the retina [14], connect to thalamic and cortical structures following a precise retinotopic ordering. The visual processing of retinal signals unfolds then through a series of stages in which low-level stimulus features are processed first by 'early' visual cortices such as V1, and increasingly complex features are processed sequentially by 'higher' extrastriate visual cortices [50].

However, feedforward connections along the visual hierarchy areas are typically complemented by reciprocal feedback connections [23, 10, 41]. For example, extra-retinal inputs to V1 may arrive from extrastriate visual areas such as V2 and V3 [54, 53, 61], but also indirectly from more distant cortical areas [66, 56, 20, 7, 15]. Moreover, reciprocal connections between V1 and visual thalamic structures have long been described, yet, their functional relevance is not yet fully understood [31, 35, 3, 69]. The integration of feedforward and feedback interactions along the visual hierarchy enables the modulation of ongoing cortical dynamics by incoming sensory input [25, 61]. In **chapter 5**, I will implement a generative model of directed interactions to examine the presence of state-dependent changes in the feedback and feedforward interactions between V1, V2 and V3.

1.2.4 Neurovascular coupling

In most of the visual cortex, blood is supplied by the calcarine and basilar arteries. Flowing through their ramifications, blood arrives in a network of small arterioles and capillary beds. After irrigating these capillary beds, which are arranged perpendicularly to the cortical surface, blood is drained through small venous channels into the dural venous sinuses. Neuroglia and astrocytes [59], the brain's housekeepers, regulate blood flow at the level of the capillary beds. These cells form part of the blood-brain barrier. They coordinate local blood

supply, nutrient transport and neurotransmitter release, and thus provide a crucial contribution to neuronal activity [12]. Thanks to this coupling between blood supply and neuronal activity, it has become possible to —indirectly— estimate neuronal population activity using non-invasive instruments like magnetic resonance scanners. In the next sections, I will show how fMRI has allowed to map the topographical and neuroanatomical organization of visual field maps in the human visual cortex.

1.2.5 Functional magnetic resonance imaging

In order to estimate neuronal activity patterns across early visual cortical areas I used fMRI. This technique measures changes in blood oxygenation, which has been shown to be indirectly related to neuronal activity [45]. Local neuronal activity induces changes in the ratio between oxyhemoglobin and deoxyhemoglobin, which can be detected due to their differential magnetic susceptibility [57]. This effect is named the blood-oxygen level dependent (BOLD) effect, which is the basis for fMRI. In my studies, I have used primarily 7T fMRI data. However, in **chapter 4**, I also examine 3T data. Higher magnetic field strengths allow for a better signal-to-noise ratio and spatiotemporal resolution. However, the temporal resolution of fMRI is limited by the hemodynamic response to neuronal activity, not by the magnetic field strength.

1.2.6 Retinotopic mapping using population receptive field (pRF) and connective field (CF) modeling

The development of functional magnetic resonance imaging (fMRI) has allowed to map the topographical and neuroanatomical organization of visual field maps in the human visual cortex [18]. Modeling the hemodynamic responses to hypothetical neuronal activity elicited by a stimulus allows the mapping of population receptive field properties [68]. The population receptive field (pRF) method relies on this approach [17]. The method uses a parameterized forward model of the neuronal population responses, a description of the hemodynamic response function (a two-gamma HRF model) [8], and the stimulus (here I used drifting checkerboard bars) to predict evoked BOLD activity. The population receptive field model used in this thesis corresponds to a circular Gaussian characterized by three parameters: x and y (positions in the stimuli screen), and size (σ).

To find the most likely retinotopic map, a set of candidate population receptive field models are combined with the stimulus aperture to generate predictions of the neuronal responses each candidate pRF would produce. Subsequent convolution of this predicted neuronal response with the HRF gives a set of candidate BOLD responses for each combination of pRF parameters. The pRF parameters associated with the best fitting candidate BOLD responses are then chosen to summarize the response of each fMRI recording site.

Somewhat similar to the way in which the visual field is mapped on the surface of the cortex using pRFs, CF modeling describes the neuronal interactions between different cortical visual areas in terms of spatial integration and cortical selectivity maps. CF modeling enables the characterization of a target record-

ing site (e.g. V2 and V3) in terms of the aggregate BOLD activity in a source brain area (e.g. V1), thus providing a description of the preferred locations on the cortical surface to which these target sites respond [32]. Locations in the primary visual cortex are associated with visual field positions obtained during visual field mapping (VFM). Therefore, visual field coordinates can be inferred for the target recording sites from the preferred locations in the source region, allowing the reconstruction of visuotopic maps even in the absence of a stimulus (e.g., RS). The technique is prominently used in **chapters 2 & 4**.

1.2.7 Resting state

Resting state (RS) research has become very popular in the last decades. Departing from the idea of the brain as a primarily reflexive device driven by the momentary demands of the environment [60, 67], RS studies suggest that the brain's operations are mainly intrinsic, involving the integration of slow internal and rapid sensory dynamics [75, 9, 44, 16]. From this perspective, sensory input modulates rather than 'forms' brain function [13].

Early evidence on the role of intrinsic brain activity came from studies of animal physiology and anatomy. Using decorticated cats, Brown [9] showed that rhythmic behaviours reminiscent of walking and running were still possible, even in the absence of sensory and cortical input. Intrinsic rhythms in brain activity were further found in humans. By recording scalp electrical potentials, Berger [5] found large-amplitude rhythmic fluctuations that appeared only when subjects had their eyes closed. These fluctuations were reported to occur at a rate of approximately 10 oscillations per second, what is now called the alpha frequency band [2]. Likewise, studying spontaneous electrical activity in the rabbit olfactory bulb, Adrian [1] found that the persistent activity of cells in the bulb unfolded as waves of electrical activity, even in the absence of sensory stimulation.

Despite these early findings, attempts to characterize and quantify intrinsic brain activity only gained momentum in the last two decades. Using multi-electrode recordings of retinal ganglion cells, Galli and Maffei [22] and Meister et al. [49] showed spontaneous wave-like propagation of neuronal firing prior to visual experience. These patterns were hypothesized to play a role in the development and formation of retinotopically ordered maps in the visual system [39, 48]. Later studies using large-scale neuronal recording approaches showed that both stimulus-evoked and spontaneous neuronal activity closely reflected the functional and anatomical organization of the visual cortex [4, 70, 40].

Parallel investigations in humans and non-human primates using fMRI recordings from the somatosensory cortex also indicated a link between the brain's intrinsic activity and its neuroanatomical organization [6, 73]. Moreover, using whole-brain fMRI recordings, it has been recently shown that the propagation of BOLD activity across distant brain areas reflects the transition between different behavioral states [51, 52]. This suggests a meaningful relationship between BOLD activity propagation and neuronal interactions between distant brain regions. Therefore, in this thesis, I ask whether task-dependent differences in BOLD activity between RS and VFM can also reveal relevant aspects of brain organization and activity, yet at a much more local scale. Thereby, I focus on

early cortical visual field maps, which are richly interconnected and for which regional variation in the hemodynamic response is less pronounced [33, 43].

1.3 References

- [1] E. D. Adrian. The electrical activity of the mammalian olfactory bulb. *Electroencephalography and Clinical Neurophysiology*, 2(1):377–388, January 1950. .
- [2] ED Adrian and BHC Matthews. The Berger Rhythm: Potential changes from the occipital lobes in man. *Brain*, 57(4):355–385, December 1934. .
- [3] Ian M Andolina, Helen E Jones, Wei Wang, and Adam M Sillito. Corticothalamic feedback enhances stimulus response precision in the visual system. *Proc. Natl. Acad. Sci. U. S. A.*, 104(5):1685–1690, 30 January 2007.
- [4] A. Arieli, D. Shoham, R. Hildesheim, and A. Grinvald. Coherent spatiotemporal patterns of ongoing activity revealed by real-time optical imaging coupled with single-unit recording in the cat visual cortex. *Journal of Neurophysiology*, 73(5):2072–2093, May 1995.
- [5] Hans Berger. Über das elektrenkephalogramm des menschen. *Archiv f. Psychiatrie*, 87(1):527–570, December 1929. .
- [6] Bharat Biswal, F Zerrin Yetkin, Victor M Haughton, and James S Hyde. Functional connectivity in the motor cortex of resting human brain using echo-planar mri. *Magn. Reson. Med.*, 34(4):537–541, 1995.
- [7] Sander E Bosch, Janneke F M Jehee, Guillén Fernández, and Christian F Doeller. Reinstatement of associative memories in early visual cortex is signaled by the hippocampus. *J. Neurosci.*, 34(22):7493–7500, 28 May 2014.
- [8] Geoffrey M. Boynton, Stephen A. Engel, Gary H. Glover, and David J. Heeger. Linear Systems Analysis of Functional Magnetic Resonance Imaging in Human V1. *Journal of Neuroscience*, 16(13):4207–4221, July 1996.
- [9] T. Graham Brown. On the nature of the fundamental activity of the nervous centres; together with an analysis of the conditioning of rhythmic activity in progression, and a theory of the evolution of function in the nervous system. *The Journal of Physiology*, 48(1):18–46, March 1914. .
- [10] J Bullier, J M Hupé, A C James, and P Girard. The role of feedback connections in shaping the responses of visual cortical neurons. *Prog. Brain Res.*, 134:193–204, 2001.
- [11] Omar H Butt, Noah C Benson, Ritobrato Datta, and Geoffrey K Aguirre. The fine-scale functional correlation of striate cortex in sighted and blind people. *J. Neurosci.*, 33(41):16209–16219, 9 October 2013.

- [12] György Buzsáki, Buzsáki György, Kaila Kai, and Raichle Marcus. Inhibition and brain work. *Neuron*, 56(5):771–783, 2007.
- [13] Felicity Callard and Daniel S Margulies. The Subject at Rest: Novel conceptualizations of self and brain from cognitive neuroscience’s study of the resting state. *Subjectivity*, 4(3):227–257, September 2011. .
- [14] Edward M. Callaway. Structure and function of parallel pathways in the primate early visual system. *The Journal of Physiology*, 566(1). .
- [15] Russell W Chan, Alex T L Leong, Leon C Ho, Patrick P Gao, Eddie C Wong, Celia M Dong, Xunda Wang, Jufang He, Ying-Shing Chan, Lee Wei Lim, and Ed X Wu. Low-frequency hippocampal-cortical activity drives brain-wide resting-state functional MRI connectivity. *Proc. Natl. Acad. Sci. U. S. A.*, 114(33):E6972–E6981, 15 August 2017.
- [16] Gustavo Deco, Morten L. Kringelbach, Viktor K. Jirsa, and Petra Ritter. The dynamics of resting fluctuations in the brain: metastability and its dynamical cortical core. *Scientific Reports*, 7(1):3095, June 2017. .
- [17] Serge O Dumoulin and Brian A Wandell. Population receptive field estimates in human visual cortex. *Neuroimage*, 39(2):647–660, 15 January 2008.
- [18] Stephen A Engel, David E Rumelhart, Brian A Wandell, Adrian T Lee, Gary H Glover, Chichilnisky Eduardo-Jose, and Michael N Shadlen. fMRI of human visual cortex. *Nature*, 369(6481):525–525, 1994.
- [19] D. J. Felleman and D. C. Van Essen. Distributed hierarchical processing in the primate cerebral cortex. *Cerebral Cortex (New York, N.Y.: 1991)*, 1(1).
- [20] Stephanie J. Forkel, Michel Thiebaut de Schotten, Jamie M. Kawadler, Flavio Dell’Acqua, Adrian Danek, and Marco Catani. The anatomy of fronto-occipital connections from early blunt dissections to contemporary tractography. *Cortex*, 56(Supplement C):73–84, July 2014. .
- [21] Karl J. Friston. Precision Psychiatry. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, 2(8):640–643, November 2017. .
- [22] Lucia Galli and Lamberto Maffei. Spontaneous impulse activity of rat retinal ganglion cells in prenatal life, October 1988.
- [23] Ralf A. W. Galuske, Kerstin E. Schmidt, Rainer Goebel, Stephen G. Lomber, and Bertram R. Payne. The role of feedback in shaping neural representations in cat visual cortex. *Proc Natl Acad Sci U S A*, 99(26):17083–17088, December 2002. .
- [24] J. J. Gibson. Ecological optics. *Vision Research*, 1(3):253–262, October 1961. .
- [25] Charles D Gilbert and Mariano Sigman. Brain states: top-down influences in sensory processing. *Neuron*, 54(5):677–696, 7 June 2007.

- [26] Matthieu Gilson, Ruben Moreno-Bote, Adrián Ponce-Alvarez, Petra Ritter, and Gustavo Deco. Estimation of directed effective connectivity from fMRI functional connectivity hints at asymmetries of cortical connectome. *PLoS Comput. Biol.*, 12(3):e1004762, March 2016.
- [27] Nicolás Gravel, Ben Harvey, Barbara Nordhjem, Koen V Haak, Serge O Dumoulin, Remco Renken, Branislava Curčić-Blake, and Frans W Cornelissen. Cortical connective field estimates from resting state fMRI activity. *Front. Neurosci.*, 8:339, 31 October 2014.
- [28] Nicolás Gravel, Ben M. Harvey, Remco J. Renken, Serge O. Dumoulin, and Frans W. Cornelissen. Phase-synchronization-based parcellation of resting state fMRI signals reveals topographically organized clusters in early visual cortex. *NeuroImage*, September 2017. .
- [29] Nicolás Gravel, Remco J. Renken, Ben M. Harvey, Gustavo Deco, Frans W. Cornelissen, and Matthieu Gilson. Propagation of BOLD activity reveals directed interactions across human visual cortex. *bioRxiv*, page 172452, October 2017. .
- [30] Amiram Grinvald, Edmund Lieke, Ron D. Frostig, Charles D. Gilbert, and Torsten N. Wiesel. Functional architecture of cortex revealed by optical imaging of intrinsic signals. *Nature*, 324(6095):361, November 1986. .
- [31] R W Guillery and S Murray Sherman. Thalamic relay functions and their role in corticocortical communication: generalizations from the visual system. *Neuron*, 33(2):163–175, 17 January 2002.
- [32] Koen V Haak, Jonathan Winawer, Ben M Harvey, Remco Renken, Serge O Dumoulin, Brian A Wandell, and Frans W Cornelissen. Connective field modeling. *Neuroimage*, 66:376–384, 1 February 2013.
- [33] Daniel A. Handwerker, John M. Ollinger, and Mark D’Esposito. Variation of BOLD hemodynamic responses across subjects and brain regions and their effects on statistical analyses. *NeuroImage*, 21(4):1639–1651, April 2004. .
- [34] Robert V Harrison, Noam Harel, Jaswinder Panesar, and Richard J Mount. Blood capillary distribution correlates with hemodynamic-based functional imaging in cerebral cortex. *Cereb. Cortex*, 12(3):225–233, March 2002.
- [35] J Michael Hasse and Farran Briggs. Corticogeniculate feedback sharpens the temporal precision and spatial resolution of visual signals in the ferret. *Proc. Natl. Acad. Sci. U. S. A.*, 114(30):E6222–E6230, 25 July 2017.
- [36] Jakob Heinzle, Thorsten Kahnt, and John-Dylan Haynes. Topographically specific functional connectivity between visual field maps in the human brain. *Neuroimage*, 56(3):1426–1436, 1 June 2011.

- [37] David H. Hubel. The Visual Cortex of the Brain. *Scientific American*, 209 (5), 1963. .
- [38] Friston Karl. A Free Energy Principle for Biological Systems. *Entropy*, 14(11):2100–2121, October 2012. .
- [39] Lawrence C. Katz and Justin C. Crowley. Development of cortical circuits: lessons from ocular dominance columns. *Nat. Rev. Neurosci.*, 3(1):34–42, January 2002. .
- [40] Tal Kenet, Dmitri Bibitchkov, Misha Tsodyks, Amiram Grinvald, and Amos Arieli. Spontaneously emerging cortical representations of visual attributes. *Nature*, 425(6961):954–956, 30 October 2003.
- [41] Bart Krekelberg. Primate motion perception. In *How Animals See the World*. Oxford University Press, 2012.
- [42] Christopher M Lewis, Conrado A Bosman, Thilo Womelsdorf, and Pascal Fries. Stimulus-induced visual cortical networks are recapitulated by spontaneous local and interareal synchronization. *Proc. Natl. Acad. Sci. U. S. A.*, 113(5):E606–15, 2 February 2016.
- [43] Fa-Hsuan Lin, Jonathan R Polimeni, Jo-Fu Lotus Lin, Kevin W-K Tsai, Ying-Hua Chu, Pu-Yeh Wu, Yi-Tien Li, Yi-Cheng Hsu, Shang-Yueh Tsai, and Wen-Jui Kuo. Relative latency and temporal variability of hemodynamic responses at the human primary visual cortex. *Neuroimage*, 22 January 2017.
- [44] R. R. Llinas. The intrinsic electrophysiological properties of mammalian neurons: insights into central nervous system function. *Science*, 242(4886): 1654–1664, December 1988. .
- [45] Nikos K Logothetis and Brian A Wandell. Interpreting the BOLD signal. *Annu. Rev. Physiol.*, 66:735–769, 2004.
- [46] Celine Mateo, Per M. Knutsen, Philbert S. Tsai, Andy Y. Shih, and David Kleinfeld. Entrainment of Arteriole Vasomotor Fluctuations by Neural Activity Is a Basis of Blood-Oxygenation-Level-Dependent Resting-State Connectivity. *Neuron*, October 2017. .
- [47] Teppei Matsui, Tomonari Murakami, and Kenichi Ohki. Transient neuronal coactivations embedded in globally propagating waves underlie resting-state functional connectivity. *Proc. Natl. Acad. Sci. U. S. A.*, 16 May 2016.
- [48] Todd McLaughlin, Christine L. Torborg, Marla B. Feller, and Dennis D. M. O’Leary. Retinotopic map refinement requires spontaneous retinal waves during a brief critical period of development. *Neuron*, 40(6):1147–1160, December 2003.
- [49] Markus Meister, Rachel O. L. Wong, Denis A. Baylor, and Carla J. Shatz. Synchronous bursts of action potentials in ganglion cells of the developing mammalian retina, May 1991.

- [50] Mortimer Mishkin, Leslie G Ungerleider, and Kathleen A Macko. Object vision and spatial vision: two cortical pathways. *Trends Neurosci.*, 6: 414–417, 1983.
- [51] Anish Mitra, Abraham Z Snyder, Enzo Tagliazucchi, Helmut Laufs, and Marcus E Raichle. Propagated infra-slow intrinsic brain activity reorganizes across wake and slow wave sleep. *Elife*, 4, 9 November 2015.
- [52] Anish Mitra, Abraham Z. Snyder, Carl D. Hacker, Mrinal Pahwa, Enzo Tagliazucchi, Helmut Laufs, Eric C. Leuthardt, and Marcus E. Raichle. Human cortical hippocampal dialogue in wake and slow-wave sleep. *PNAS*, 113(44):E6868–E6876, November 2016. .
- [53] Lars Muckli and Lucy S Petro. Network interactions: non-geniculate input to V1. *Curr. Opin. Neurobiol.*, 23(2):195–201, April 2013.
- [54] S Murray Sherman and R W Guillery. *Functional Connections of Cortical Areas: A New View from the Thalamus*. MIT Press, 9 August 2013.
- [55] Ian Nauhaus and Kristina J. Nielsen. Building maps from maps in primary visual cortex. *Curr. Opin. Neurobiol.*, 24(1):1–6, February 2014. .
- [56] Behrad Noudoost and Tirin Moore. Control of visual cortical signals by prefrontal dopamine. *Nature*, 474(7351):372–375, 15 May 2011.
- [57] S Ogawa, T M Lee, A R Kay, and D W Tank. Brain magnetic resonance imaging with contrast dependent on blood oxygenation. *Proc Natl Acad Sci U S A*, 87(24):9868–9872, December 1990.
- [58] Pratik Y O’Herron, Philip andand Chhatbar, Levy Manuel, Shen Zhiming, Adrien E Schramm, Lu Zhongyang, and Kara Prakash. Neural correlates of single-vessel haemodynamic responses in vivo. *Nature*, 2016.
- [59] J C Pang, P A Robinson, K M Aquino, and N Vasan. Effects of astrocytic dynamics on spatiotemporal hemodynamics: Modeling and enhanced data analysis. *Neuroimage*, 14 October 2016.
- [60] I. P. Pavlov and For Mem R. S. Croonian Lecture. Certain problems in the physiology of the cerebral hemispheres. *Proc. R. Soc. Lond. B*, 103 (722):97–110, June 1928. .
- [61] Lucy S Petro, Luca Vizioli, and Lars Muckli. Contributions of cortical feedback to sensory processing in primary visual cortex. *Front. Psychol.*, 5: 1223, 6 November 2014.
- [62] Mathijs Raemaekers, Raemaekers Mathijs, Schellekens Wouter, Richard J A van Wezel, Petridou Natalia, Kristo Gert, and Nick F Ramsey. Patterns of resting state connectivity in human primary visual cortical areas: A 7T fMRI study. *Neuroimage*, 84:911–921, 2014.
- [63] M E Raichle. The restless brain: how intrinsic activity organizes brain function. *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, 370(1668):20140172–20140172, 2015.

- [64] Lars Reichl, Dominik Heide, Siegrid Löwel, Justin C. Crowley, Matthias Kaschube, and Fred Wolf. Coordinated Optimization of Visual Cortical Maps (I) Symmetry-based Analysis. *PLoS Computational Biology*, 8(11), 2012. .
- [65] Mark M Schira, Christopher W Tyler, Michael Breakspear, and Branka Spehar. The foveal confluence in human visual cortex. *J. Neurosci.*, 29(28): 9050–9058, 15 July 2009.
- [66] Jeremy D. Schmahmann and Deepak N. Pandya. The Complex History of the Fronto-Occipital Fasciculus. *Journal of the History of the Neurosciences*, 16(4):362–377, October 2007. .
- [67] Charles Sherrington. Ferrier Lecture: Some Functional Problems Attaching to Convergence. *Proceedings of the Royal Society of London. Series B, Containing Papers of a Biological Character*, 105(737):332–362, 1929.
- [68] A T Smith, K D Singh, A L Williams, and M W Greenlee. Estimating receptive field size from fMRI data in human striate and extrastriate visual cortex. *Cereb. Cortex*, 11(12):1182–1190, December 2001.
- [69] M. Steriade. Corticothalamic resonance, states of vigilance and mentation. *Neuroscience*, 101(2):243–276, 2000.
- [70] M. Tsodyks, T. Kenet, A. Grinvald, and A. Arieli. Linking Spontaneous Activity of Single Cortical Neurons and the Underlying Functional Architecture. *Science*, 286(5446):1943–1946, December 1999. .
- [71] Francisco J Varela. Autopoiesis and a Biology of Intentionality. 1991.
- [72] Alberto L Vazquez, Matthew C Murphy, and Seong-Gi Kim. Neuronal and physiological correlation to hemodynamic resting-state fluctuations in health and disease. *Brain Connect.*, 4(9):727–740, November 2014.
- [73] J L Vincent, G H Patel, M D Fox, A Z Snyder, J T Baker, D C Van Essen, J M Zempel, L H Snyder, M Corbetta, and M E Raichle. Intrinsic functional architecture in the anaesthetized monkey brain. *Nature*, 447(7140):83–86, 3 May 2007.
- [74] Brian A Wandell, Serge O Dumoulin, and Alyssa A Brewer. Visual field maps in human cortex. *Neuron*, 56(2):366–383, 25 October 2007.
- [75] James William. The Principles of Psychology. *Philosophy and Phenomenological Research*, 44(1):124–126, 1983.